

Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <a href="https://www.bccancer.bc.ca/terms-of-use">www.bccancer.bc.ca/terms-of-use</a> and according to acceptable standards of care.

PROTOCOL CODE: BRAJACTW

Page 1 of 2

DOCTOR'S ORDERS Htcm Wtkg BSA	m²				
REMINDER: Please ensure drug allergies and previous bleomycin are documented on the Allergy & Alert Form					
DATE: To be given: Cycle #:					
Date of Previous Cycle:					
Delay treatment week(s)  CBC & Diff, platelets day of treatment  May proceed with doses as written for AC portion if labs done within 96 h: ANC greater than or element  Platelets greater than or equal to 90 x 109/L  May proceed with doses as written for weekly paclitaxel portion if labs done within 24 h: ANC greater than or equal to 90 x 109/L, Platelets greater than or equal to 90 x 109/L  Dose modification for: Hematology Other Toxicity  Proceed with treatment based on blood work from	eater than or equal to 1.0				
PREMEDICATIONS: Patient to take own supply. RN/Pharmacist to confirm					
dexamethasone    8 mg or    12 mg (select one) PO 30 to 60 minutes prior to AC treatment and select ONE of the following:  ondansetron 8 mg PO 30 to 60 minutes prior to AC treatment aprepitant 125 mg PO 30 to 60 minutes prior to AC treatment ondansetron 8 mg PO 30 to 60 minutes prior to AC treatment netupitant-palonosetron 300 mg-0.5 mg PO 30 to 60 minutes prior to AC treatment					
45 Minutes Prior to PACLitaxel: <b>dexamethasone 10 mg</b> IV in 50 mL NS over 15 minutes 30 Minutes Prior to PACLitaxel: <b>diphenhydrAMINE 25 mg</b> IV in NS 50 mL over 15 minutes and NS 100 mL over 15 minutes (Y-site compatible)  No pre-medication required (see protocol for guidelines)  Other:					
**Have Hypersensitivity Reaction Tray and Protocol Available for Cycles	5 to 8**				
CHEMOTHERAPY:  DOXOrubicin 60 mg/m² x BSA =mg  Dose Modification:% =mg/m² x BSA =mg  IV push  cyclophosphamide 600 mg/m² x BSA =mg  Dose Modification:% =mg/m² x BSA =mg  IV in 100 to 250 mL NS over 20 minutes to 1 hour  OR					
PACLitaxel  80 mg/m² OR  mg/m² (select one) x BSA = mg  Dose Modification:  % = mg/m² x BSA = mg  IV in 100 to 500 mL (non-DEHP bag) NS over 1 hour once weekly x 3 weeks (use non-DEHP line filter)  DOSE MODIFICATION IF REQUIRED ON SUBSEQUENT DAYS (Cycle 5-8):	tubing with 0.2 micron in-				
PACLitaxel 80 mg/m² x BSA = mg  Dose Modification: % = mg/m² x BSA = mg  IV in 100 to 500 mL (non-DEHP bag) NS over 1 hour on days (use non-DEH in-line filter)	IP tubing with 0.2 micron				
DOCTOR'S SIGNATURE:	SIGNATURE: UC:				



Information on this form is a guide only. User will be solely responsible for verifying its currency and accuracy with the corresponding BC Cancer treatment protocols located at <a href="https://www.bccancer.bc.ca/terms-of-use">www.bccancer.bc.ca/terms-of-use</a> and according to acceptable standards of care.

PROTOCOL CODE: BRAJACTW

Page 2 of 2

DOCTOR'S ORDERS	Ht	cm Wt	kg	BSA	m²	
DATE:						
RETURN APPOINTMENT ORDERS						
Return in three weeks for Doctor and Cycle for cycles 5-8)	(Bo	ok chemo room weekly	/ x 3			
Last Cycle. Return inweek	κ(s) after last tr	eatment.		l		
CBC & Diff, Platelets prior to each treatment  Prior to Cycle #5: Bilirubin, ALT  If clinically indicated: Creatinine ALT Bilir  Other tests:  Consults:  See general orders sheet for additional reque		A	1			
DOCTOR'S SIGNATURE:				SIGNAT UC:	URE:	