

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Leucovorin 50 mg/5 mL 200 mg/20 mL 1000 mg/100 mL (GMP) (F)(PFL) no preservative ¹	N/A	10 mg/mL ¹	50 mg: discard unused portion ^{1,2} 200 mg, 1000 mg: 8 h F ^{1,2}	syringe	8 h RT ^{1,2}	
				0.05-10 mg/mL NS, D5W, Ringer's, LR, D10W, D5-NS ^{1,2} 50-250 mL†	NS , D5W, LR, Ringer's: 24 h RT ¹ D10W, D5-NS: 8 h RT ¹	
Leucovorin 50 mg/5 mL 500 mg/50 mL (Pfizer/Hospira) (F)(PFL) no preservative ³	N/A	10 mg/mL ³	8 h ³	syringe	8 h RT ³	
				0.05–10 mg/mL NS , D5W, LR, Ringer's, D10W, D5NS ³ 50-250 mL†	NS , D5W, LR, Ringer's: 24 h RT ³ D10W, D5NS: 8 h RT ³	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Leucovorin 50 mg/5 mL 500 mg/50 mL (Teva) (F)(PFL) no preservative ⁴	N/A	10 mg/mL ⁵	discard unused portion ⁵	syringe	8 h ^{6,7}	
				0.4 - 4.8 mg/mL NS , D5W ⁸ 50-250 mL†	72 h F, RT ⁸	
				0.06 - 0.4 mg/mL NS , D5W ⁴ 50-250 mL†	NS: 24 h RT ⁴ D5W: 12 h RT ⁴	
				0.06 - 1 mg/mL Ringer's, Lactated Ringer's, D10W, D10-NS ⁴	Ringer's, LR: 24 h RT ⁴ D10W: 12 h RT ⁴ D10NS: 6 h RT ⁴	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Lurbinectedin 4 mg (Jazz) (F) no preservative ⁹	8 mL SWI ⁹	0.5 mg/mL ⁹	12 h F , RT ^{9,10}	100-250 mL NS , D5W ⁹	complete administration within 24 h F , RT ⁹	- larger infusion volume is recommended for peripheral line ⁹ - do not use nylon membrane filters for administration if diluted in NS ⁹ ; BD Alaris pumps and syringe sets have polyethersulfone membrane in-line filters ¹¹
Lurbinectedin 4 mg (Pharma Mar) (F) no preservative ¹² (SAP)	8 mL SWI ¹²	0.5 mg/mL ¹²	12 h F , RT ^{10,12}	100–250 mL NS , D5W ¹²	30 h F , RT ¹²	- larger infusion volume is recommended for peripheral line ¹²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Melphalan 50 mg (Marcan) (RT)(PFL) no preservative ¹³	10 mL supplied diluent ¹³ rapidly add diluent and immediately shake vigorously to dissolve ¹³ record time of reconstitution	5 mg/mL ¹³	2 h RT ¹³ do NOT refrigerate¹³	0.1-0.45 mg/mL NS only¹³	complete administration within 50 min RT from time of initial reconstitution ¹³	- will precipitate if stored in fridge ¹³
Melphalan 50 mg (Taro) (RT)(PFL) no preservative ¹⁴	10 mL supplied diluent ¹⁴ rapidly add diluent and immediately shake vigorously to dissolve ¹⁴ record time of reconstitution	5 mg/mL ¹⁴	2 h RT ¹⁴ do NOT refrigerate¹⁴	0.1-0.45 mg/mL NS only¹⁴	complete administration within 50 min RT from time of initial reconstitution ¹⁴	- will precipitate if stored in fridge ¹⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mesna 400 mg/4 mL 1000 mg/10 mL (Baxter) (RT) no preservative ¹⁵	N/A	100 mg/mL ¹⁵ (use filter needle to withdraw from ampoule)	discard unused portion ¹⁵	≥1 mg/mL NS , D5W, D5½-NS, LR ¹⁵⁻¹⁷ 100 mL†	24 h RT ¹⁵	
Mesna 1000 mg/10 mL 5000 mg/50 mL (Baxter) (RT) preservative ¹⁵	N/A	100 mg/mL ¹⁵	8 d RT ¹⁵ (vial may be punctured up to 4 times) ¹⁵	≥1 mg/mL NS , D5W, D5½-NS, LR ¹⁵⁻¹⁷ 100 mL†	24 h RT ¹⁵	
Mesna 1000 mg/10 mL (Fresenius Kabi) (RT) preservative ¹⁸	N/A	100 mg/mL ¹⁸	14 d F , RT ^{18,19}	≥1 mg/mL NS , D5W ²⁰ 100 mL†	48 h F, 24 h RT ¹⁸	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Methotrexate 50 mg/2 mL 500 mg/20 mL 1 g/40 mL (Accord) (RT)(PFL) no preservative ²¹	N/A	25 mg/mL ²¹	50mg: discard unused portion ²¹ 500 mg, 1 g: 8 h RT ²¹	syringe	use within 8 h RT of initial puncture ²¹	- for high-dose regimens (e.g., 1-12 g/m ² as a single dose): use preservative-free methotrexate ²¹ - do not use for IT injection
				0.4–2 mg/mL NS , D5W ²¹ 50-500 mL†	use within 24 h RT of initial puncture ²¹ **(PFL)	
				high dose (e.g., 1-12 g/m ² as a single dose): 1000 mL * NS	use within 24 h RT of initial puncture ²¹ **(PFL)	
Methotrexate intravitreal injection 50 mg/2 mL (Accord) (RT)(PFL) no preservative ²¹	N/A	25 mg/mL ²¹	discard unused portion ²¹	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	- for intravitreal use preservative-free methotrexate is preferred ²²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Methotrexate IT Injection Only preservative free methotrexate may be administered by the intrathecal route²³ 50 mg/2 mL (Accord) (RT)(PFL) no preservative²¹</p>	N/A	25 mg/mL ²¹	discard unused portion ²¹	<p>IT syringe</p> <p>qs to 6 mL with preservative free NS^{24,25}</p> <p>diluents containing preservatives should NOT be used for intrathecal administration²⁶</p>	use within 4 h of initial puncture ¹⁰	<p>- auxiliary info¹⁰: IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer Ziploc bag²⁷</p>
<p>Methotrexate 50 mg/2 mL 500 mg/20 mL (Accord) (RT)(PFL) preservative²¹</p>	N/A	25 mg/mL ²¹	28 d F ^{10,21}	<p>syringe</p> <p>0.4–2 mg/mL NS, D5W²¹</p> <p>50-500 mL†</p>	<p>10 d F^{10,21}</p> <p>24 h RT²¹</p>	<p>- contains benzyl alcohol²¹ - do NOT use for high-dose regimens (e.g., 1-12 g/m² as a single dose)²¹ - do NOT use for IT injection²¹</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Methotrexate 50 mg/2 mL 500 mg/20 mL 1 g/40 mL 2.5 g/100 mL (Pfizer/Hospira) (RT)(PFL) no preservative ²⁸	N/A	25 mg/mL ²⁸	50mg: discard unused portion ²⁸ 500 mg, 1 g, or 2.5 g: 8 h RT ²⁸	syringe	use within 8 h RT of initial puncture ²⁸	- for high-dose regimens (e.g., 1-12 g/m ² as a single dose): use preservative-free methotrexate ²⁸ - do not use for IT injection
				0.4–2 mg/mL NS , D5W ²⁸ 50-500 mL†	use within 24 h RT of initial puncture ²⁸ **(PFL)	
				high dose (e.g., 1-12 g/m ² as a single dose): 1000 mL * NS	use within 24 h RT of initial puncture ²⁸ **(PFL)	
Methotrexate intravitreal injection 50 mg/2 mL (Pfizer/Hospira) (RT)(PFL) no preservative ²⁸	N/A	25 mg/mL ²⁸	discard unused portion ²⁸	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	- for intravitreal use preservative-free methotrexate is preferred ²²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Methotrexate IT Injection Only preservative free methotrexate may be administered by the intrathecal route ²³ 50 mg/2 mL (Pfizer/Hospira) (RT)(PFL) no preservative ²⁸	N/A	25 mg/mL ²⁸	discard unused portion ²⁸	IT syringe qs to 6 mL with preservative free NS ^{24,25} diluents containing preservatives should NOT be used for intrathecal administration ²⁶	use within 4 h of initial puncture ¹⁰	- auxiliary info ¹⁰ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer Ziploc bag ²⁷
Methotrexate 50 mg/2 mL 500 mg/20 mL (Pfizer/Hospira) (RT)(PFL) preservative ²⁸	N/A	25 mg/mL ²⁸	28 d F ^{10,28}	syringe	10 d F ^{10,28}	- contains benzyl alcohol ²⁸ - do NOT use for high-dose regimens (e.g., 1-12 g/m ² as a single dose) ²⁸ - do NOT use for IT injection ²⁸
				0.4–2 mg/mL NS, D5W ²⁸ 50-500 mL†	24 h RT ²⁸	
Mitomycin 20 mg (Accord) (RT)(PFL) no preservative ²⁹	40 mL SWI ²⁹ shake well ²⁹	0.5 mg/mL ²⁹	12 h F, 6 h RT ^{10,30} **(PFL) ³⁰	syringe	72 h F, 6 h RT ³⁰ **(PFL) ³⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mitomycin intravesical 20 mg (Accord) (RT)(PFL) no preservative ²⁹	40 mL SWI ²⁹ shake well ²⁹	0.5 mg/mL ²⁹	12 h F, 6 h RT ^{10,30} **(PFL) ³⁰	syringe	72 h F, 6 h RT ³⁰ **(PFL) ³⁰	
	10 mL SWI ³¹ shake well ²⁹	2 mg/mL ³¹	use immediately after preparation to prevent precipitation ³²	syringe	use immediately after preparation to prevent precipitation ³²	- may precipitate due to low solubility ^{32,33} - do NOT refrigerate ³²
	25 mL SWI shake well	0.8 mg/mL ³⁴	discard unused portion ^{2,34} **(PFL) ^{2,34}	syringe	4 days RT ³⁴ **(PFL) ^{2,34}	- do NOT refrigerate ³⁴
	33.3 mL SWI shake well	0.6 mg/mL ³⁴	discard unused portion ^{2,34} **(PFL) ^{2,34}	syringe	4 days F, RT ³⁴ **(PFL) ^{2,34}	
Mitomycin intraperitoneal 20 mg (Accord) (RT)(PFL) no preservative ²⁹	40 mL SWI ²⁹ shake well ²⁹	0.5 mg/mL ²⁹	12 h F, 6 h RT ^{10,30} **(PFL) ³⁰	0.02-0.04 mg/mL NS , sodium lactate ²⁹	NS: 18 h F, 3 h RT ³⁰ sodium lactate: 6 h F, 3 h RT ³⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mitomycin 20 mg (Teva) (RT)(PFL) no preservative ³⁵	40 mL SWI ³⁵ shake well ³⁵	0.5 mg/mL ³⁵	12 h F, 6 h RT ^{10,35} **(PFL) ³⁵	syringe	72 h F, 6 h RT ³⁵ **(PFL) ³⁵	
Mitomycin intravesical 20 mg (Teva) (RT)(PFL) no preservative ³⁵	40 mL SWI ³⁵ shake well ³⁵	0.5 mg/mL ³⁵	12 h F, 6 h RT ^{10,35} **(PFL) ³⁵	syringe	72 h F, 6 h RT ³⁵ **(PFL) ³⁵	
	10 mL SWI ³¹ shake well ³⁵	2 mg/mL ³¹	use immediately after preparation to prevent precipitation ³²	syringe	use immediately after preparation to prevent precipitation ³²	- may precipitate due to low solubility ^{32,33} - do NOT refrigerate ³²
	25 mL SWI shake well	0.8 mg/mL ³⁴	discard unused portion ^{2,34} **(PFL) ^{2,34}	syringe	4 days RT ³⁴ **(PFL) ^{2,34}	- do NOT refrigerate ³⁴
	33.3 mL SWI shake well	0.6 mg/mL ³⁴	discard unused portion ^{2,34} **(PFL) ^{2,34}	syringe	4 days F, RT ³⁴ **(PFL) ^{2,34}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mitomycin intraperitoneal 20 mg (Teva) (RT)(PFL) no preservative ³⁵	40 mL SWI ³⁵ shake well ³⁵	0.5 mg/mL ³⁵	12 h F, 6 h RT ^{10,35} **(PFL) ³⁵	0.02-0.04 mg/mL NS , sodium lactate ³⁵	NS: 18 h F, 6 h RT ³⁵ sodium lactate: 6 h F, RT ³⁵	
mitoXANTRONE 20 mg/10 mL (Fresenius Kabi) (RT) no preservative ³⁶	N/A	2 mg/mL ³⁶	discard unused portion ³⁶	0.2-0.6 mg/mL NS , D5W ³⁶ 50 mL†	24 h RT ³⁶	
mitoXANTRONE 20 mg/10 mL 25 mg/12.5 mL 30 mg/15 mL (Pfizer/Hospira) (RT)(PFL) no preservative ³⁷	N/A	2 mg/mL ³⁷	discard unused portion ³⁷	0.2-0.6 mg/mL NS , D5W ³⁷ 50 mL†	72 h F, 24 h RT ³⁷ **(PFL) ³⁷	
Mogamulizumab 20 mg/5 mL (Kyowa) (F)(PFL) do not shake no preservative ³⁸	N/A	4 mg/mL ³⁸	discard unused portion ³⁸	0.1-3 mg/mL NS 100 mL* mix by gentle inversion; do not shake ³⁸	24 h F ³⁸	- discard if cloudy, discoloured, or visible particulates are present ³⁸ - administer with 0.2 micron in-line filter ³⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Mogamulizumab 20 mg/5 mL (Kyowa) (F)(PFL) do not shake no preservative ³⁹ (SAP)	N/A	4 mg/mL ³⁹	discard unused portion ³⁹	0.1-3 mg/mL NS ³⁹ 100 mL* mix by gentle inversion; do not shake ³⁹	24 h F ³⁹	- discard if cloudy or discoloured ³⁹ - administer with 0.2 micron in-line filter ³⁹
Nivolumab 40 mg/4 mL 100 mg/10 mL (BMS) (F)(PFL) do not shake no preservative ⁴⁰	N/A	10 mg/mL ⁴⁰	discard unused portion ⁴⁰	1-10 mg/mL NS , D5W ⁴⁰ 25-100 mL† mix by gentle inversion; do not shake ⁴⁰ OR undiluted in empty infusion bag or glass bottle ⁴⁰	complete administration within 7 days F, including max 8 h at RT ⁴⁰ **(PFL) ⁴⁰ (can be in room light when at RT) ⁴⁰	- do not shake ⁴⁰ - administer with 0.2 micron in-line filter ⁴⁰ - may contain a few amorphous particles ⁴⁰ - discard if cloudy, has pronounced colour change (should be clear to pale yellow) ⁴⁰

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
oBINutuzumab 1000 mg/40 mL (Roche) (F)(PFL)** do not shake no preservative ⁴¹	N/A	25 mg/mL ⁴¹	discard unused portion ⁴²	NS 100 mg: 100 mL ⁴¹ 900 mg: 250 mL ⁴¹ 1000 mg: 250 mL ⁴¹	24 h F, 48 h RT ^{41,43}	-once removed from the fridge, diluted product is stable for an additional 48 h RT ^{41,43} - do NOT shake ⁴¹ - do NOT use dextrose containing solutions ⁴¹
Octreotide 50 mcg/1 mL 100 mcg/1 mL 500 mcg/1 mL (Omega) (F)(PFL) no preservative ⁴⁴	N/A	50 mcg/mL ⁴⁴	discard unused portion ⁴⁴	NS ⁴⁴ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/h ⁴⁴	24 h RT ⁴⁴	
		100 mcg/mL ⁴⁴				
		500 mcg/mL ⁴⁴				
Octreotide multidose vial: 1000 mcg/5 mL (Omega) (F)(PFL) preservative ⁴⁴	N/A	200 mcg/mL ⁴⁴	15 d F ⁴⁴	NS ⁴⁴ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/h ⁴⁴	24 h RT ⁴⁴	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Octreotide (SANDOSTATIN®) 50 mcg/1 mL 100 mcg/1 mL 500 mcg/1 mL (Novartis) (F)(PFL) no preservative ⁴⁵	N/A	50 mcg/mL ⁴⁵	discard unused portion ⁴⁵	NS ⁴⁵ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/h ⁴⁵	24 h RT ⁴⁵	
		100 mcg/mL ⁴⁵				
		500 mcg/mL ⁴⁵				
Octreotide (SANDOSTATIN®) multi-dose vial: 1000 mcg/5 mL (Novartis) (F)(PFL) preservative ⁴⁵	N/A	200 mcg/mL ⁴⁵	14 d F, RT ⁴⁵	NS ⁴⁵ volume adjusted to ensure a continuous infusion of octreotide at 25 mcg/h ⁴⁵	24 h RT ⁴⁵	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Octreotide (SANDOSTATIN LAR®) (long acting) 10 mg 20 mg 30 mg (Novartis) (F)(PFL) no preservative⁴⁵</p>	<p>2 mL supplied diluent⁴⁵</p> <p>add diluent: gently run diluent down sides of vial⁴⁵</p> <p>do NOT disturb for 2–5 min; then swirl moderately⁴⁵</p> <p>record time of reconstitution</p>	<p>10 mg: 5 mg/mL⁴⁵</p> <hr/> <p>20 mg: 10 mg/mL⁴⁵</p> <hr/> <p>30 mg: 15 mg/mL⁴⁵</p>	<p>discard unused portion⁴⁵</p>	<p>syringe (for deep intragluteal administration only)⁴⁵</p>	<p>use within 4 h of initial reconstitution^{10,45}</p>	<p>- do NOT shake⁴⁵</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Octreotide suspension (long acting) 10 mg 20 mg 30 mg (Teva) (F)(PFL) no preservative⁴⁶</p>	2 mL supplied diluent	10 mg: 5 mg/mL ⁴⁶	discard unused portion ⁴⁶	syringe (for deep intragluteal administration only) ⁴⁶	use within 4 h of initial reconstitution ^{10,46}	- gently shake to resuspend before administration ⁴⁶ - delay in administration may result in sedimentation ⁴⁶
	let stand at RT for 30 min prior to reconstitution ⁴⁶	20 mg: 10 mg/mL ⁴⁶				
	<p>add supplied diluent⁴⁶</p> <p>let vial stand for 5 min after adding diluent to saturate powder⁴⁶</p> <p>shake moderately in horizontal direction for ≥30 sec to create suspension⁴⁶</p> <p>record time of reconstitution</p>	30 mg: 15 mg/mL ⁴⁶				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Dr. Reddy's) (RT)(PFL) no preservative ⁴⁷	N/A	5 mg/mL ⁴⁷	discard unused portion ⁴⁷	0.2-0.7 mg/mL D5W ⁴⁷ 100-500 mL† do NOT use NS or other chloride- containing solution ⁴⁷ do NOT use aluminum-containing needle and syringe ⁴⁷	0.2-2 mg/mL: 48 h F, 24 h RT ⁴⁷	- do NOT use aluminum- containing needle, syringe, or tubing ⁴⁷
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Pfizer/Hospira) (RT) no preservative ⁴⁸	N/A	5 mg/mL ⁴⁸	discard unused portion ⁴⁸	0.2-0.7 mg/mL D5W ⁴⁸ 100-500 mL† do NOT use NS or other chloride- containing solutions ⁴⁸ do NOT use aluminum-containing needle and syringe ⁴⁸	0.2-0.4 mg/mL: 24 h RT ⁴⁸ or 5 d F plus an additional 8 h RT ⁴⁹ 0.5–2 mg/mL: 24 h RT ⁴⁸ or 10 d F, plus an additional 8 h RT ^{10,49} **(PFL) when stored in F ⁴⁹	- do NOT use aluminum- containing needle, syringe, tubing ⁴⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 150 mg/30 mL 200 mg/40 mL (Sandoz) (RT)(PFL) no preservative ⁵⁰	N/A	5 mg/mL ⁵⁰	12 h F, RT ^{10,51}	0.2-0.7 mg/mL D5W ⁵⁰ 100-500 mL† do NOT use NS or other chloride- containing solution ⁵⁰ do NOT use aluminum-containing needle and syringe ⁵⁰	0.2-2 mg/mL: 48 h F, 24 h RT ⁵⁰	- do NOT use aluminum- containing needle, syringe, tubing ⁵⁰
Oxaliplatin 50 mg/10 mL 100 mg/20 mL 200 mg/40 mL (Teva) (RT)(PFL) no preservative ⁵²	N/A	5 mg/mL ⁵²	discard unused portion ⁵²	0.2-0.7 mg/mL D5W ⁵² 100-500 mL† do NOT use NS or other chloride- containing solution ⁵² do NOT use aluminum-containing needle and syringe ⁵²	0.2-2 mg/mL: 48 h F, 24 h RT ⁵²	- do NOT use aluminum- containing needle, syringe or tubing ⁵²

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
PACLitaxel 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Accord) (RT)(PFL) preservative ⁵³	N/A	6 mg/mL ⁵³	30 mg, 100 mg: 28 d RT ^{10,53} 300 mg: 24 h RT ^{10,53}	0.3-1.2 mg/mL NS , D5W, D5NS, D5LR ⁵³ 50-500 mL†	complete administration within 27 h RT ⁵³	- use non-DEHP bag and tubing ⁵³ - administer with 0.2 micron in-line filter ⁵³ - avoid excessive shaking ⁵³
				0.1 mg/mL NS ⁵⁴	44 h F , RT ⁵⁴	
PACLitaxel 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Biolyse) (RT) preservative ⁵⁵	N/A	6 mg/mL ⁵⁵	28 d RT ⁵⁶	0.3-1.2 mg/mL NS , D5W ⁵⁵ 50-500 mL†	complete administration within 27 h RT ^{57,58}	- use non-DEHP bag and tubing ⁵⁵ - administer with 0.2 micron in-line filter ⁵⁵
				0.1 mg/mL NS ⁵⁴	44 h F , RT ⁵⁴	
				0.012-0.12 mg/mL NS ⁵⁹	16 h RT ⁵⁷	
				devices with spikes (e.g., chemo dispensing pins) may be used with vials ⁶⁰		

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Paclitaxel 30 mg/5 mL 100 mg/16.7 mL 300 mg/50 mL (Sandoz) (RT)(PFL) preservative ⁶¹	N/A	6 mg/mL ⁶¹	30 mg, 100 mg: 28 d RT ^{10,61} 300 mg: 24 h RT ^{10,61}	0.3-1.2 mg/mL NS , D5W, D5NS ⁶¹ 50-500 mL†*	complete administration within 27 h RT ⁶¹	- use non-DEHP bag and tubing ⁶¹ - administer with 0.2 micron inline filter ⁶¹ - avoid excessive shaking
				0.1 mg/mL NS ⁵⁴	44 h F , RT ⁵⁴	
PACLitaxel, nanoparticle, albumin- bound (NAB) 100 mg (Celgene) (RT)(PFL) no preservative ⁶²	20 mL NS ⁶² slowly direct diluent against side of vial (i.e., ≥1 min) during reconstitution ⁶² let stand for ≥5 min to wet powder ⁶² gently swirl or invert for ≥2 min ⁶²	5 mg/mL ⁶²	use immediately (RT) or 8 h F ⁶² **(PFL) ⁶²	in empty sterile PVC, non-PVC, or non- DEHP infusion bag ⁶²	48 h F plus an additional 8 h RT ⁶³	- each vial contains 900 mg human albumin ⁶² - to prevent foaming, do NOT inject NS directly onto the powder ⁶² - some settling may occur; use mild agitation to resuspend ⁶² - administer with 15 micron filter ONLY ⁶² (NOTE: filters with pore size less than 15 microns may cause filter blockage) ⁶⁴

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
PACLitaxel, nanoparticle, albumin- bound (NAB) 100 mg (Panacea/Apo) (RT)(PFL) no preservative ⁶⁵	20 mL NS ⁶⁵ slowly direct diluent against side of vial (i.e., ≥1 min) during reconstitution ⁶⁵ let stand for ≥5 min to wet powder ⁶⁵ gently swirl or invert for ≥2 min ⁶⁵ (if foaming occurs, let stand for ≥15 min) ⁶⁵	5 mg/mL ⁶⁵	use immediately (RT) or 8 h F ⁶⁵ **(PFL) ⁶⁵	in empty sterile PVC, non-PVC, or non-DEHP infusion bag ⁶⁵	56 h F plus an additional 4 h RT ⁶⁶	<ul style="list-style-type: none"> - each vial contains 900 mg human albumin⁶⁵ - to prevent foaming, do NOT inject NS directly onto the powder⁶⁵ - some settling may occur; use gentle inversion to resuspend⁶⁵ - discard if visible particulates are present⁶⁵ - administer with 15 micron filter ONLY⁶⁵
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Fresenius Kabi) (RT) no preservative ⁶⁷	N/A	3 mg/mL ⁶⁷	discard unused portion ⁶⁷	≤0.36 mg/mL ⁶⁷ NS , D5W ⁶⁷ 250 mL†	24 h RT ⁶⁷	<ul style="list-style-type: none"> - do NOT mix with calcium containing solutions (e.g., Lactated Ringer's)⁶⁷
		6 mg/mL ⁶⁷				
		9 mg/mL ⁶⁷				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Hospira) (RT) no preservative ⁶⁸	N/A	3 mg/mL ⁶⁸	discard unused portion ⁶⁸	0.06–0.36 mg/mL NS , D5W ⁶⁸ 250 mL†	24 h F plus an additional 24 h RT (total 48 h) ⁶⁸ **(PFL) ⁶⁸	- do NOT mix with calcium containing solution (e.g., Lactated Ringer's) ⁶⁸
		6 mg/mL ⁶⁸				
		9 mg/mL ⁶⁸				
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Omega) (RT) no preservative ⁶⁹	N/A ⁶⁹	3 mg/mL ⁶⁹	discard unused portion ⁶⁹	0.06–0.36 mg/mL NS , D5W ⁶⁹ 250 mL†	24 h F plus an additional 24 h RT (total 48 h) ⁶⁹ **(PFL) ⁶⁹	- do NOT mix with calcium containing solution (e.g., Lactated Ringer's) ⁶⁹
		6 mg/mL ⁶⁹				
		9 mg/mL ⁶⁹				
Pamidronate 30 mg/10 mL 60 mg/10 mL 90 mg/10 mL (Pfizer) (RT) no preservative ⁷⁰	N/A	3 mg/mL ⁷⁰	discard unused portion ⁷⁰	0.06-0.36 mg/mL NS , D5W ⁷⁰ 250 mL†	24 h F plus an additional 24 h RT (total 48 h) ⁷⁰ **(PFL) ⁷⁰	- do NOT mix with calcium containing solution (e.g., Lactated Ringer's) ⁷⁰
		6 mg/mL ⁷⁰				
		9 mg/mL ⁷⁰				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pamidronate 30 mg/10 mL 60mg/10 mL 90 mg/10 mL (Sandoz Canada) RT no preservative ⁷¹	N/A	3 mg/mL ⁷¹	discard unused portion ^{71,72}	NS ; D5W ⁷¹ 250 mL†	24 h RT ⁷¹	- do NOT mix with calcium containing solution (e.g., Lactated Ringer's) ⁷¹
		6 mg/mL ⁷¹				
		9 mg/mL ⁷¹				
PANitumumab 100 mg/5 mL 400 mg/20 mL (Amgen) (F)(PFL) do not shake no preservative ⁷³	N/A	20 mg/mL ⁷³	discard unused portion ⁷³	1-10mg/mL NS ⁷³ 100 mL†	24 h F, 6 h RT ⁷³⁻⁷⁶	- administer with 0.2 micron in-line filter ⁷³ - solution may contain particulates which do not affect product quality ⁷³ - do not administer if discoloured ⁷³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pegaspargase (pegylated asparaginase <i>E. coli</i>) 3750 units/5 mL (Servier) (F)(PFL) do not shake no preservative ⁷⁷	N/A	750 units/mL ⁷⁷	discard unused portion ⁷⁷	IM ⁷⁷ : max volume: 2 mL in children and adolescents; 3 mL in adults if volume greater than above, use multiple sites ⁷⁷	syringe: use within 4 h of vial puncture ^{2,77}	- do NOT shake ⁷⁷
				IV ⁷⁷ : 100 mL NS , D5W	bag: use within 4 h of vial puncture ^{2,77}	
Pembrolizumab 100 mg/4 mL (Merck) (F)(PFL) do not shake no preservatives ⁷⁸	N/A	25 mg/mL ⁷⁸	discard unused portion ^{2,78}	1-10 mg/mL NS , D5W ⁷⁸ 50 mL * mix by gentle inversion ⁷⁸	complete administration within 96 h F, 6 h RT ⁷⁸	- administer with 0.2 micron in-line filter ⁷⁸ - bring vials and diluted solutions to RT prior to use ⁷⁸ - vials contain 0.25 mL overflow ⁷⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pemetrexed 100 mg 500 mg (Accord) (RT) no preservative ⁷⁹	100 mg: 4.2 mL NS ⁷⁹ 500 mg: 20 mL NS ⁷⁹	25 mg/mL ⁷⁹	12 h F, RT ^{10,79}	100 mL NS ⁷⁹	24 h F, RT ⁷⁹	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁷⁹
Pemetrexed 100 mg/4 mL 500 mg/20 mL 850 mg/34 mL 1000 mg/40 mL (Accord) (RT)(PFL) no preservative ⁸⁰	N/A	25 mg/mL ⁸⁰	discard unused portion ⁸⁰	100 mL NS ⁸⁰	24 h F ⁸⁰	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁸⁰
Pemetrexed 100 mg 500 mg (Dr. Reddy's) (RT) no preservative ⁸¹	100 mg: 4.2 mL NS ⁸¹ 500 mg: 20 mL NS ⁸¹	25 mg/mL ⁸¹	12 h F, RT ^{10,82-84}	100 mL NS ⁸¹	24 h F, RT ⁸²⁻⁸⁴	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁸¹
Pemetrexed 100 mg 500 mg (Lilly) (RT) no preservative ⁸⁵	100 mg: 4.2 mL NS ⁸⁵ 500 mg: 20 mL NS ⁸⁵	25 mg/mL ⁸⁵	12 h F ^{10,85}	100 mL NS ⁸⁵	24 h F ⁸⁵	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁸⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Pemetrexed 100 mg 500 mg 1000 mg (Taro) (RT) no preservative ⁸⁶	100 mg: 4.2 mL NS ⁸⁶ 500 mg: 20 mL NS ⁸⁶ 1000 mg: 40 mL NS ⁸⁶	25 mg/mL ⁸⁶	12 h F ^{10,86}	100 mL NS ⁸⁶	24 h F ⁸⁶	- do NOT mix with calcium containing solution (e.g., Ringer's) ⁸⁶
Pentostatin 10 mg (Hospira/Pfizer) (F) no preservative ⁸⁷	5 mL SWI ⁸⁷	2 mg/mL ⁸⁷	8 h RT ⁸⁷	0.18-0.33 mg/mL ⁸⁷ 25-50 mL NS, D5W ⁸⁷	8 h RT ⁸⁷	
PERTuzumab 420 mg/14 mL (Roche) (F)(PFL) no preservative ⁸⁸	N/A	30 mg/mL ⁸⁸ do NOT shake ⁸⁸	discard unused portion ^{42,88}	250 mL NS only ⁸⁸ mix by gentle inversion to avoid foaming ⁸⁸	24 h F, RT ⁸⁸	- do NOT use dextrose containing solutions ⁸⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>PERTuzumab- trastuzumab 1200 mg-600 mg/15 mL 600 mg-600 mg/10 mL (Roche) (F)(PFL) do not shake no preservative⁸⁹</p>	<p>N/A</p>	<p>1200 mg-600 mg⁸⁹: 80 mg/mL pertuzumab and 40 mg/mL trastuzumab</p> <p>600 mg-600 mg⁸⁹: 60 mg/mL pertuzumab and 60 mg/mL trastuzumab</p>	<p>discard unused portion⁸⁹</p>	<p>SC syringe⁸⁹</p>	<p>10 d F, 24 h RT^{10,89}</p>	<p>- do not shake⁸⁹ - contains recombinant human hyaluronidase⁸⁹</p>
<p>Plerixafor 24 mg/1.2 mL (sanofi-aventis) (RT) no preservative⁹⁰</p>	<p>N/A</p>	<p>20 mg/mL⁹⁰</p>	<p>discard unused portion⁹⁰</p>	<p>SC syringe⁹⁰</p>	<p>48 h RT^{72,91}</p>	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Polatuzumab vedotin 30 mg 140 mg (Hoffman-La Roche) (F)(PFL) do not shake no preservative ⁹²	30 mg: 1.8 mL SWI ⁹² 140 mg: 7.2 mL SWI ⁹² direct diluent against side of vial during reconstitution ⁹² swirl gently to mix ⁹²	20 mg/mL ⁹² (PFL)	12 h F, RT ^{10,92}	0.72-2.7 mg/mL NS , D5W, ½NS ⁹² (dilute to a minimum volume of 50 mL) ⁹² gently invert bag to mix ⁹²	in NS: 72 h F, 4 h RT ⁹² in D5W or ½NS: 72 h F, 8 h RT ⁹²	- do NOT shake ⁹² - administer with 0.2 micron in-line filter ⁹² -discard if discolouration or visible particulates are present ⁹²
Pralatrexate 20 mg/1 mL 40 mg/2 mL (Servier) (F)(PFL) no preservative ⁹³	N/A	20 mg/mL ⁹³	discard unused portion ²	syringe ⁹³	24 h F, RT ⁹⁴ **(PFL) ⁹⁴	- do NOT dilute ⁹³
Raltitrexed 2 mg (Pfizer) (F,RT)(PFL) no preservative ⁹⁵	4 mL SWI ⁹⁵	0.5 mg/mL ⁹⁵	12 h F, RT ^{10,95}	50-250 mL NS , D5W ⁹⁵	complete administration within 24 h F, RT ⁹⁵	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Ramucirumab 100 mg/10 mL 500 mg/50 mL (Eli Lilly) (F)(PFL) (do not shake) no preservative ⁹⁶	N/A	10 mg/mL ⁹⁶	discard unused portion ⁹⁶	0.4–4 mg/mL NS ^{96,97} 250-500 mL† gently invert to mix ⁹⁶ do NOT shake ⁹⁶	24 h F, 4 h RT ⁹⁶	- administer with 0.2 micron in-line filter ⁹⁶ - do NOT use dextrose containing solutions ⁹⁶
riTUXimab (RITUXAN®) 100 mg/10 mL 500 mg/50 mL (Roche) (F)(PFL) no preservative ⁹⁸	N/A	10 mg/mL ⁹⁸	discard unused portion ⁹⁸	1-4 mg/mL NS, D5W ⁹⁸ 250-500 mL†	NS: 10 d F plus an additional 24 h RT ^{10,98} D5W: 24 h F plus an additional 12 h RT ⁹⁸	
riTUXimab intravitreal injection (RITUXAN®) 100 mg/10 mL (Roche) (F)(PFL) no preservative ⁹⁸	N/A	10 mg/mL ⁹⁸	discard unused portion ⁹⁸	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
riTUXimab subcutaneous (RITUXAN® SC) 1400 mg/11.7 mL 1600 mg/13.4 mL (Roche) (F)(PFL) no preservative ⁹⁹	N/A	120 mg/mL ⁹⁹	discard unused portion ⁹⁹	SC syringe ⁹⁹	48 h F plus 8 h RT ⁹⁹	- contains hyaluronidase ⁹⁹ - formulations are NOT interchangeable ⁹⁹
riTUXimab (RIXIMYO®) 100 mg/10 mL 500 mg/50 mL (Sandoz) (F)(PFL) (do NOT shake) no preservative ¹⁰⁰	N/A	10 mg/mL ¹⁰⁰	discard unused portion ¹⁰⁰	1-4 mg/mL NS, D5W ¹⁰⁰ 250-500 mL† gently invert to mix	NS: 10 d F plus an additional 24 h RT ^{10,100} D5W: 24 h F plus an additional 12 h RT ¹⁰⁰	
riTUXimab intravitreal injection (RIXIMYO®) 100 mg/10 mL (Sandoz) (F)(PFL) (do NOT shake) no preservative ¹⁰⁰	N/A	10 mg/mL ¹⁰⁰	discard unused portion ¹⁰⁰	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
riTUXimab (RUXIENCE®) 100 mg/10 mL 500 mg/50 mL (Pfizer) (F)(PFL) no preservative ¹⁰¹	N/A	10 mg/mL ¹⁰¹	discard unused portion ¹⁰¹	1-4 mg/mL NS , D5W ¹⁰¹ 250-500 mL† gently invert to mix	24 h F plus an additional 24 h RT ¹⁰¹	
riTUXimab intravitreal injection (RUXIENCE®) 100 mg/10 mL 500 mg/50 mL (Pfizer) (F)(PFL) no preservative ¹⁰¹	N/A	10 mg/mL ¹⁰¹	discard unused portion ¹⁰¹	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	
riTUXimab (TRUXIMA®) 100 mg/10 mL 500 mg/50 mL (Teva/Celltrion) (F)(PFL) no preservative ¹⁰²	N/A	10 mg/mL ¹⁰²	discard unused portion ¹⁰²	1-4 mg/mL NS , D5W ¹⁰² 250-500 mL† gently invert to mix	24 h F plus an additional 12 h RT ¹⁰²	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
riTUXimab intravitreal injection (TRUXIMA®) 100 mg/10 mL 500 mg/50 mL (Teva/Celltrion) (F)(PFL) no preservative ¹⁰²	N/A	10 mg/mL ¹⁰²	discard unused portion ¹⁰²	syringe for intravitreal use	use within 4 h of initial puncture ¹⁰	
romiDEPsin 10 mg (Celgene Inc.) (RT) ¹⁰³ no preservative ⁴²	2.2 mL supplied diluent ^{103,104} swirl gently to mix ¹⁰³	5 mg/mL ¹⁰³	8 h RT ¹⁰³	500 mL NS ¹⁰³	24 h RT ¹⁰³	- reconstituted solution will be slightly viscous ¹⁰⁵ - vials contain overflow to allow for full drug recovery (drug vial contains 11 mg romidepsin; diluent vial contains 2.4 mL diluent) ¹⁰³

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Sacituzumab govitecan 180 mg (Gilead) (F)(PFL) no preservative ¹⁰⁶	20 mL NS ¹⁰⁶ bring vials to RT before reconstitution ¹⁰⁶ slowly add diluent to vial and gently swirl; allow to dissolve for up to 15 min ¹⁰⁶ do not shake ¹⁰⁶	10 mg/mL ¹⁰⁶	use immediately after reconstitution to prepare infusion solution ¹⁰⁶ discard unused portion ¹⁰⁶	1.1-3.4 mg/mL NS ¹⁰⁶ 100-1000 mL NS† slowly inject solution to bag to minimize foaming; do not shake ¹⁰⁶	24 h F ¹⁰⁶ , plus an additional 8 h RT including infusion time ¹⁰⁶ **(PFL) ¹⁰⁶	- do not shake ¹⁰⁶ - protect container from light during administration ¹⁰⁶ - vials contain overflow (~20 mg per vial) ¹⁰⁷
Siltuximab 100 mg 400 mg (Janssen) (F)(PFL) no preservative ¹⁰⁸	100 mg: 5.2 mL SWI ¹⁰⁸ 400 mg: 20 mL SWI ¹⁰⁸ bring vial to RT prior to use (~30 min) ¹⁰⁸ gently swirl, do NOT shake ¹⁰⁸	20 mg/mL ¹⁰⁸	2 h RT ¹⁰⁸	250 mL D5W ¹⁰⁸ dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ¹⁰⁸	complete administration within 6 h RT ¹⁰⁸	- administer with 0.2 micron in-line filter ¹⁰⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Sirolimus, nanoparticle, albumin- bound (NAB) 100 mg (Aadi) (F)(PFL) no preservative¹⁰⁹ (SAP)</p>	<p>20 mL NS¹⁰⁹ slowly direct diluent against side of vial (over ≥1 min)¹⁰⁹ let stand for ≥5 min to wet powder¹⁰⁹ gently swirl or invert for ≥2 min to avoid foaming¹⁰⁹ if foaming/clumping occurs, let stand until foam subsides (≥15 min)¹⁰⁹</p>	<p>5 mg/mL¹⁰⁹</p>	<p>4 h F^{110,111} **(PFL)¹⁰⁹</p>	<p>undiluted in empty PVC or non-PVC infusion bag¹⁰⁹</p>	<p>9 h F, followed by max 4 h RT¹⁰⁹ **(PFL)¹⁰⁹</p>	<p>- each vial contains ~800-900 mg human albumin^{109,112} - to prevent foaming, do NOT inject NS directly onto the powder¹⁰⁹ - if powder is visible after reconstitution, gently invert to resuspend powder¹⁰⁹ - to prevent administration of proteinaceous strands, administer with 15 micron filter ONLY¹⁰⁹</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Streptozocin 1g (Keocyt) (F)(PFL) no preservative ¹¹³⁻¹¹⁶ (SAP)	9.5mL NS , SWI, D5W ¹¹³⁻¹¹⁶	100 mg/mL ¹¹³⁻¹¹⁶	12 h F ^{10,114-116}	syringe ¹¹⁴⁻¹¹⁶	48 h F ^{10,114-116}	
				100-500 mL NS , D5W, SWI ¹¹³⁻¹¹⁶	24 h F ¹¹⁴⁻¹¹⁶	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Tebentafusp 100 mcg/0.5 mL (Immunocore/Medison) (F)(PFL) do not shake no preservative¹¹⁷</p>	<p>N/A</p>	<p>200 mcg/mL¹¹⁷</p>	<p>discard unused portion¹¹⁷</p>	<p>100 mL NS¹¹⁷</p> <p>Step 1: add calculated volume of human albumin 5% to provide 225-275 mcg/mL final concentration¹¹⁷</p> <p>to mix: invert the bag and gently rotate ≥5 times; do NOT shake bag (repeat x3)¹¹⁷</p> <p>Step 2: add calculated volume of drug¹¹⁷</p> <p>to mix: invert the bag and gently rotate ≥5 times; do NOT shake bag (repeat x3)¹¹⁷</p>	<p>complete administration within 24 h F, 4 h RT¹¹⁷</p> <p>bring to RT prior to administration¹¹⁷</p>	<p>- do NOT use CSTD or filters during preparation¹¹⁷ - CSTD can be used for administration¹¹⁸ - administer using 0.2 micron in-line filter¹¹⁷ - once the bag has been removed from fridge, it must remain at RT¹¹⁷</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Tebentafusp 100 mcg/0.5 mL (Immunocore/Clinigen) (F)(PFL) do not shake no preservative^{119,120} (SAP)</p>	<p>N/A</p>	<p>200 mcg/mL¹¹⁹</p>	<p>discard unused portion^{2,119,120}</p>	<p>100 mL NS^{119,120}</p> <p>Step 1: add calculated volume of human albumin 5% to provide 225-275 mcg/mL final concentration^{119,120}</p> <p>to mix: invert the bag and gently rotate ≥5 times; do NOT shake bag (repeat x3)^{119,120}</p> <p>Step 2: add calculated volume of drug^{119,120}</p> <p>to mix: invert the bag and gently rotate ≥5 times; do NOT shake bag (repeat x3)^{119,120}</p>	<p>complete administration within 24 h F, 4 h RT^{119,120}</p>	<ul style="list-style-type: none"> - do NOT use CSTD or filters during preparation¹¹⁹ - CSTD can be used for administration¹¹⁸ - administer using 0.2 micron in-line filter^{119,120} - once the bag has been removed from fridge, it must remain at RT^{119,120}

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Teclistamab 30 mg/3 mL 153 mg/1.7 mL (Janssen) (F)(PFL) do not shake no preservative¹²¹</p>	<p>N/A</p>	<p>30 mg¹²¹: 10 mg/mL</p> <p>(use for 2.1-52.9 mg doses)*</p>	<p>discard unused portion¹²¹</p>	<p>SC syringe¹²¹</p> <p>if drug volume >2 mL, divide volume into separate syringes for administration¹²¹</p>	<p>20 h F, RT¹²¹</p> <p>if stored in fridge, bring to RT prior to administration¹²¹</p>	<p>- CAUTION: two concentrations are available¹²¹ - CSTD must not be used for preparation or administration of syringe volumes less than 1 mL¹²²; use filtered venting needle (e.g., Chemo-Vent®) in place of CSTD for preparation¹²³</p>
		<p>153 mg¹²¹: 90 mg/mL</p> <p>(use for 53-375 mg doses)*</p>				
		<p>bring to RT before use (~15 min)¹²¹</p> <p>swirl gently for 10 sec to mix; do NOT shake¹²¹</p>				

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Temsirolimus 30 mg/1.2 mL (Pfizer/Wyeth) (F)(PFL) ¹²⁴ no preservative ¹²⁵	1.8 mL supplied diluent ¹²⁴	10 mg/mL ¹²⁴	12 h RT ^{10,124} **(PFL) ¹²⁴	250 mL NS ¹²⁴ record time of dilution ¹²⁴	complete administration within 6 h ¹²⁴ mix by gentle inversion to avoid foaming ¹²⁴	- use non-DEHP bag and tubing - administer with 0.2 micron in-line filter ¹²⁴
Teniposide 50 mg/5 mL (BMS) (RT) preservative ¹²⁶	N/A	10 mg/mL ¹²⁶	discard unused portion	0.1-1 mg/mL NS , D5W ¹²⁶ 50–500 mL *	0.1-0.4 mg/mL: 24 h RT ¹²⁶ 1 mg/mL: complete administration within 4 h RT of preparation ^{126,127}	- do not refrigerate - use non-DEHP bag and tubing ¹²⁶ - do not use if precipitates ^{126,127} - contains DMA*** - excessive agitation may cause precipitation ¹²⁶

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Thiotepa 15 mg 100 mg (Adienne/Methapharm) (F) no preservative ¹²⁸ (SAP)	15 mg: 1.5 mL SWI ¹²⁸ 100 mg: 10 mL SWI ¹²⁸ to remove haze, filter through 0.22 micron filter after reconstitution ¹²⁹ record time of reconstitution	10 mg/mL ¹²⁸	8 h F ¹²⁸	0.5-1 mg/mL NS ¹²⁸ ≤500 mg: 500 mL ¹²⁸ >500 mg: 1000 mL ¹²⁸ reconstituted solution is hypotonic and must be further diluted with NS prior to use ¹²⁸	24 h F, 4 h RT ¹²⁸	- do not use if precipitates are present ¹²⁸ - reconstituted solution may be used if opalescent ¹²⁸ - administer with 0.2 micron in-line filter ¹²⁸

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Thiotepa IT injection 15 mg 100mg (Adienne/Methapharm) (F) no preservative ¹²⁸ (SAP)	15 mg: 1.5 mL SWI ¹²⁸ 100 mg: 10 mL SWI ¹²⁸ diluents containing preservatives should NOT be used for intrathecal administration ²⁶ to remove haze, filter through 0.22 micron filter after reconstitution ¹²⁹ record time of reconstitution	10 mg/mL ¹²⁸	8 h F ¹²⁸	IT syringe qs to 6 mL with preservative free NS ¹³⁰ diluents containing preservatives should NOT be used for intrathecal administration ²⁶	use within 4 h of initial reconstitution ²	- auxiliary info ²⁷ : IT - label to include route in full (i.e., INTRATHECAL injection) attached to both syringe and outer ziplock bag ²⁷ - do not use if precipitates are present ¹²⁸ - reconstituted solution may be used if opalescent ¹²⁸
Thyrotropin alfa 1.1 mg (Genzyme) (F)(PFL) no preservative ¹³¹	1.2 mL SWI ¹³¹ swirl gently to mix ¹³¹ do NOT shake ¹³¹	0.9 mg/mL ¹³¹	12 h F ^{10,131}	syringe ¹³¹	24 h F ^{10,131}	- do not use if particulates are present ¹³¹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Tislelizumab 100 mg/10 mL (BeiGene) (F) ^{132,133} (do not shake) no preservative ¹³⁴ (SAP)	N/A	10 mg/mL ¹³⁴	discard unused portion ¹³⁴	1-10 mg/mL NS ¹³⁴ 50-100 mL*	complete administration within 20 h F, 4 h RT (max 24 h from preparation) ¹³⁴ bring to RT prior to administration ¹³⁴ mix by gentle inversion; do not shake ¹³⁴	
Tocilizumab 80 mg/4 mL 200 mg/10 mL 400 mg/20 mL (Roche) (F)(PFL) no preservative ¹³⁵	N/A	20 mg/mL ¹³⁵	discard unused portion ¹³⁵	100 mL NS ¹³⁵ dilute to final volume by withdrawing volume from bag equal to volume of drug to be added ¹³⁵ gently invert to mix ¹³⁵	complete administration within 24 h F, RT ¹³⁵ bring to RT prior to administration ¹³⁵	- to prevent foaming: slowly add drug to infusion bag and gently invert bag to mix ¹³⁵
Topotecan 4 mg/4 mL (Accord) (RT)(PFL) no preservative ¹³⁶	N/A	1 mg/mL ¹³⁶	12 h F, RT ^{10,136}	0.025-0.5 mg/mL NS, D5W ¹³⁶ 25-50 mL†	10 d F, 4 d RT ^{10,136}	

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Topotecan 4 mg/4 mL (Pfizer/Hospira) (F)(PFL) no preservative ¹³⁷	N/A	1 mg/mL ¹³⁷	discard unused portion ¹³⁷	0.02-0.5 mg/mL NS , D5W ¹³⁷ 25-50 mL†	24 h F, RT ¹³⁷	
Topotecan 4 mg/4 mL (Sandoz) (F)(PFL) no preservative ¹³⁸	N/A	1 mg/mL ¹³⁸	discard unused portion ¹³⁸	0.02-0.5 mg/mL NS , D5W ¹³⁸ 25-50 mL†	24 h F ¹³⁸ **(PFL) ¹³⁸	
Trastuzumab (HERCEPTIN®) 440 mg (Roche) (F) no preservative ¹³⁹	20 mL supplied BWI ¹³⁹ swirl vial gently; allow to stand undisturbed for 5 min ¹³⁹	21 mg/mL ¹³⁹	28 d F ¹³⁹	250 mL NS only ¹³⁹ do NOT use dextrose containing solutions ¹³⁹	24 h F, RT ¹³⁹	- do NOT shake ¹³⁹

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Trastuzumab (HERZUMA®) 150 mg 440 mg (Teva/Celltrion) (F) no preservative ¹⁴⁰	150 mg: 7.2 mL SWI ¹⁴⁰	21 mg/mL ¹⁴⁰	discard unused portion ¹⁴⁰	250 mL NS only ¹⁴⁰ do NOT use dextrose containing solutions ¹⁴⁰	24 h F, RT ¹⁴⁰	- do NOT shake ¹⁴⁰ - supplied BWI contains benzyl alcohol ¹⁴⁰
	440 mg: 20 mL supplied BWI ¹⁴⁰		28 d F ¹⁴⁰			
	swirl vial gently; allow to stand undisturbed for 5 min ¹⁴⁰					
Trastuzumab (OGIVRI®) 150 mg 440 mg (BGP) (F) no preservative ¹⁴¹	150 mg: 7.2 mL SWI ¹⁴¹	21 mg/mL ¹⁴¹	discard unused portion ¹⁴¹	250 mL NS only ¹⁴¹ do NOT use dextrose containing solutions ¹⁴¹	24 h F, RT ¹⁴¹	- do NOT shake ¹⁴¹ - supplied BWI contains benzyl alcohol ¹⁴¹
	440 mg: 20 mL supplied BWI ¹⁴¹		28 d F ¹⁴¹			
	swirl vial gently; allow to stand undisturbed for 5 min ¹⁴¹					

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Trastuzumab (TRAZIMERA®) 150 mg 440 mg (Pfizer) (F) no preservative ¹⁴²	150 mg: 7.2 mL SWI ¹⁴²	21 mg/mL ¹⁴²	discard unused portion ¹⁴²	250 mL NS only ¹⁴² do NOT use dextrose containing solutions ¹⁴²	24 h F, RT ¹⁴²	- do NOT shake ¹⁴² - supplied BWI contains benzyl alcohol ¹⁴²
	440 mg: 20 mL supplied BWI ¹⁴²		28 d F ¹⁴²			
	swirl vial gently; allow to stand undisturbed for 5 min ¹⁴²					

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Trastuzumab deruxtecan (ENHERTU®) 100 mg (AstraZeneca) (F)(PFL) no preservative¹⁴³</p>	<p>5 mL SWI¹⁴³ swirl gently until completely dissolved¹⁴³ do NOT shake¹⁴³</p>	<p>20 mg/mL¹⁴³</p>	<p>12 h F^{10,143} **(PFL)¹⁴³</p>	<p>100 mL D5W only¹⁴³ gently invert to mix¹⁴³ do NOT shake¹⁴³ do NOT use sodium chloride solution¹⁴³</p>	<p>complete administration within 24 h F, 4 h RT¹⁴³ **(PFL)¹⁴³</p>	<p>- do not use if reconstituted solution contains visible particulates or is cloudy or discoloured¹⁴³ - protect container from light during administration¹⁴⁴ - administer with 0.2 micron in-line filter¹⁴³ - if stored in fridge, bring bag to RT prior to use¹⁴³</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Trastuzumab emtansine (KADCYLA®) 100 mg 160 mg (Roche) (F)(PFL) no preservative¹⁴⁵</p>	<p>100 mg: 5 mL SWI¹⁴⁵</p> <p>160 mg: 8 mL SWI¹⁴⁵</p> <p>swirl gently until completely dissolved</p> <p>do NOT shake¹⁴⁵</p>	<p>20 mg/mL¹⁴⁵</p>	<p>12 h F^{10,146}</p>	<p>250 mL NS or ½NS only¹⁴⁵</p> <p>do NOT shake¹⁴⁵</p> <p>do NOT use dextrose containing solutions¹⁴⁵</p>	<p>24 h F¹⁴⁵</p>	<p>- do not use if reconstituted solution contains visible particulates or is cloudy or discolored¹⁴⁵ - D5W causes aggregation of the protein¹⁴⁵ - for infusions prepared in NS: administer with 0.2 micron in-line filter or 0.22 micron polyethersulfane (PES) filter¹⁴⁵ - for infusions prepared in ½NS: filter is optional for administration¹⁴⁵</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
<p>Treosulfan 1 g 5 g (Medexus) (RT) no preservative¹⁴⁷</p>	<p>1 g¹⁴⁷: 20 mL NS, D5W, SWI, ½NS</p> <p>5 g¹⁴⁷: 100 mL NS, D5W, SWI, ½NS</p> <p>pre-heat diluent to 25-30°C (max)¹⁴⁸</p> <p>shake vial to loosen powder before adding the warmed diluent¹⁴⁹</p> <p>vigorous shaking may be required¹⁴⁹; prolonged standing time may improve solubility¹⁴⁷</p>	<p>50 mg/mL¹⁴⁷</p>	<p>12 h RT^{10,147}</p>	<p>undiluted in empty infusion bag^{147,148}</p>	<p>3 d RT¹⁴⁷</p>	<p>- do NOT refrigerate as may precipitate¹⁴⁷</p>

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Treosulfan 1 g 5 g (medac) (RT) no preservative ^{150,151} (SAP)	1 g ^{150,151} : 20 mL SWI, ½NS 5 g ^{150,151} : 100 mL SWI, ½NS pre-heat diluent to 25-30°C (max) ^{150,151} shake vial carefully to loosen powder before adding the warmed diluent ^{150,151} gently shake while adding diluent ^{150,151} (takes ~2 min to reconstitute) ^{150,151}	50 mg/mL ^{150,151}	12 h RT ^{10,150,152}	undiluted ¹⁵³ or dilute with NS or D5W in empty infusion bag for final concentration = 20 mg/mL ¹⁵²	4 d RT ^{150,152}	- compatible with polytetrafluoroethyl ene filters ¹⁵⁴ - may sometimes require vigorous shaking to reconstitute ^{150,151} - do NOT refrigerate as may cause precipitation ^{150,151}
vinBLAStine 10 mg/10 mL (Pfizer) (F)(PFL) no preservative ¹⁵⁵	N/A	1 mg/mL ¹⁵⁵	discard unused portion ^{2,155}	25-50 mL NS , D5W ¹⁵⁶	use within 4 h of initial vial puncture ^{2,155}	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES^{157,158}

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
vinBLAS tine 10 mg/10 mL (Teva) (F)(PFL) no preservative ¹⁵⁹	N/A	1 mg/mL ¹⁵⁹	discard unused portion ^{2,159}	25-50 mL NS , D5W ¹⁵⁶	use within 4 h of initial vial puncture ^{2,159}	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{157,158}
vinCRIS tine 2 mg/2 mL 5 mg/5 mL (Pfizer/Hospira) (F)(PFL) no preservative ¹⁶⁰	N/A	1 mg/mL ¹⁶⁰	8 h F, RT ¹⁶⁰	0.01-0.1 mg/mL NS , D5W ¹⁶⁰ 50 mL†	24 h F, RT ¹⁶⁰ **(PFL) ¹⁶⁰	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{157,158} - for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRIStine)

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
vinCRISTine 1 mg/1 mL 2 mg/2 mL 5 mg/5 mL (Teva) (F)(PFL) no preservative ¹⁶¹	N/A	1 mg/mL ¹⁶¹	8 h F, RT ¹⁶¹	0.01-0.1 mg/mL NS , D5W ¹⁶¹ 50 mL†	24 h F, RT ¹⁶¹	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES^{157,158} - for LYEPOCHR protocol, see entry for EPOCHR (3-in-1 solution containing either etoposide or etoposide phosphate AND DOXOrubicin and vinCRISTine)
Vinorelbine 10 mg/1 mL 50 mg/5mL (Fresenius Kabi) (F)(PFL) no preservative ¹⁶²	N/A	10 mg/mL ¹⁶²	discard unused portion ¹⁶²	0.5-2.0 mg/mL NS , D5W, ½NS, D5-½NS, Ringer's, Ringer's Lactate ¹⁶² 50 mL†	24 h F, RT ¹⁶²	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES^{157,158}

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Vinorelbine 10 mg/1 mL 50 mg/5 mL (GMP) (F)(PFL) no preservative ¹⁶³	N/A	10 mg/mL ¹⁶³	discard unused portion ²	0.5-2.0 mg/mL NS , D5W, ½NS, D5-½NS, Ringer's, Ringer's Lactate ¹⁶³ 50 mL†	24 h F, RT ¹⁶³	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{157,158}
Vinorelbine 10 mg/1 mL 50 mg/5 mL (Teva) (F)(PFL) no preservative ¹⁶⁴	N/A	10 mg/mL ¹⁶⁴	discard unused portion ¹⁶⁴	0.5–2.0 mg/mL NS , D5W, ½NS, D5-½NS, Ringer's, Ringer's Lactate ¹⁶⁴ 50 mL†	24 h F, RT ¹⁶⁴	- auxiliary info: WARNING: FOR INTRAVENOUS USE ONLY – FATAL IF GIVEN BY OTHER ROUTES ^{157,158}
Zoledronic acid 4 mg/5 mL (Dr Reddy's) (RT) no preservative ¹⁶⁵	N/A	0.8 mg/mL ¹⁶⁵	discard unused portion ¹⁶⁵	100 mL NS , D5W ¹⁶⁵	complete infusion within 24 h of preparation ¹⁶⁵ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁶⁵	- do NOT mix with calcium containing solutions (e.g., Lactated Ringer's) ¹⁶⁵

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Zoledronic acid 4 mg/5 mL (Marcan) (RT) no preservative ¹⁶⁶	N/A	0.8 mg/mL ¹⁶⁶	discard unused portion ¹⁶⁶	100 mL NS , D5W ¹⁶⁶	complete infusion within 24 h of preparation ¹⁶⁶ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁶⁶	- do NOT mix with calcium containing solutions (e.g., Lactated Ringer's) ¹⁶⁶
Zoledronic acid 4 mg/5 mL (MDA) (RT) no preservative ¹⁶⁷	N/A	0.8 mg/mL ¹⁶⁷	discard unused portion ¹⁶⁷	100 mL NS , D5W ¹⁶⁷	complete infusion within 24 h of preparation ¹⁶⁷ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁶⁷	- do NOT mix with calcium containing solutions (e.g., Lactated Ringer's) ¹⁶⁷

BC CANCER CHEMOTHERAPY PREPARATION AND STABILITY CHART

DRUG & STRENGTH (Storage Prior to Use, Manufacturer, Preservative Status)	Reconstitute With:	To Give:	Vial Stability	Product (for IV bag size selection, see Notes†)	Product Stability	Special Precautions/Notes
Zoledronic acid (ZOMETA) 4 mg/ 5 mL (Novartis) (RT) no preservative ¹⁶⁸	N/A	0.8 mg/mL ¹⁶⁸	discard unused portion ⁴²	100 mL NS, D5W ¹⁶⁸	complete infusion within 24 h of preparation ¹⁶⁸ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁶⁸	- do NOT mix with calcium containing solutions (e.g., Lactated Ringer's) ¹⁶⁸
Zoledronic acid 4 mg/5 mL (Sandoz) (RT) no preservative ¹⁶⁹	N/A	0.8 mg/mL ¹⁶⁹	discard unused portion ¹⁶⁹	100 ml NS, D5W ¹⁶⁹	complete infusion within 24 h of preparation ¹⁶⁹ refrigerate diluted product if not used immediately after preparation; bring to RT prior to use ¹⁶⁹	- do NOT mix with calcium containing solutions (e.g., Lactated Ringer's) ¹⁶⁹

* Suggested volume based on usual dose range and any concentration range of stability data

† see [BC Cancer IV Bag Selection table](#): standardized bag sizes are provided for select Benefit Drugs with concentration-dependent stability or large drug volume

** Protect from light means minimizing exposure to direct sunlight over a *storage* period. More specific information on protection from light (eg, protecting container and tubing during *administration*) will be indicated in the Special Precautions/Notes column.

*** Contains DMA (N,N dimethylacetamide). Product may be incompatible with closed system transfer devices (CSTD) such as ChemoLock.

Centres are not to change content locally. All suggestions for change are to be forwarded to the Cancer Drug Manual editor.

Explanatory Notes:

Stability data assumes products prepared using standard aseptic technique in biological safety cabinet at low risk for contamination according to the classification outlined in USP 797.^{170,171}

Vial stability: Stability of solution after first puncture or reconstituted solution.

Storage temperature: If information states same stability with refrigerator and room temperature storage, then fridge stability is bolded as preferred (ie, to minimize growth of micro-organisms).

Discard unused portion: Unused portion from single use vials should be discarded at the end of the day.

“overflow known” is stated if the manufacturer states overflow that is present is within acceptable limits.

“Complete administration within ___” is stated if the manufacturer specifies that the infusion must be completed in a specific time frame following preparation, usually including entire time required for preparation (from first puncture), storage, and administration of infusion.

Nomenclature for **In-line filters** has been standardized to 0.2 micron filter size. For more information, refer to CDM monograph.

Abbreviations:

BWI = bacteriostatic water for injection

CIVI: ambulatory pump = Continuous Intravenous Infusion (e.g., elastomeric infusor)

CSTD = closed system transfer device

D5W = dextrose 5% in water

DMA = N,N dimethylacetamide

F = refrigerate

Non-DEHP = not containing Di(2-ethylhexyl) phthalate (DEHP)

NS = normal saline

PFL = protect from light

RT = room temperature

SAP = drug is approved for use through the Health Canada Special Access Program

SWI = sterile water for injection

References:

1. Generic Medical Partners Inc. Leucovorin calcium injection product monograph. Toronto, Ontario; 6 March 2020
2. BC Cancer. Provincial Pharmacy Directive Number II-20: Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; December 5 2018
3. Pfizer Canada Inc. Leucovorin calcium injection product monograph. Kirkland, Quebec; 21 June 2018
4. Teva Canada Limited. Leucovorin calcium injection® product monograph. Toronto, Ontario; 5 May 2014
5. Hospira Healthcare Corporation. LEUCOVORIN CALCIUM INJECTION® product monograph. Saint-Laurent, Quebec; 7 June . 2007
6. Novopharm Limited (Teva). LEUCOVORIN CALCIUM® Injection product information package. Toronto, Ontario; undated undated
7. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 6 January 2006
8. Jenny Yeung. Medical Information Specialist, Teva Canada. Personal communication. 12 April 2017

9. Jazz Pharmaceuticals Canada Incorporated. ZEPZELCA® product monograph. Mississauga, Ontario; September 29, 2021
10. BC Cancer. Provincial Pharmacy Directive Number II-20: Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; May 1 2022
11. Linda Ewing, BD US Market Manager. WW Infusion Disposables. Personal communication. Feb 2018
12. Pharma Mar. Lurbinectedin Compassionate Use: Preparation Guide for Infusion - edition 1. Madrid, Spain; April 2019
13. Marcan Pharmaceuticals Inc. Melphalan for injection product monograph. Ottawa, ON; July 26, 2019
14. Taro Pharmaceuticals Inc. Taro-Melphalan product monograph. Brampton, Ontario; April 5, 2019
15. Baxter Corporation. UROMITEXAN® product monograph. Mississauga, Ontario; 6 August 2013
16. Mona Ghobros BPharm MSc. Medical Information, Baxter Corporation. Personal communication. 29 November 2018
17. Trissel's® 2 Clinical Pharmaceutics Database (database on the Internet). Mesna. Lexi-Comp Inc.; created by Lawrence A. Trissel, Available at: <http://online.lexi.com>. Accessed 29 November, 2018
18. Fresenius Kabi Canada Ltd. Mesna for injection product monograph. Richmond Hill, Ontario; 21 December 2017
19. BC Cancer. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer; 19 September 2007
20. Fresenius Kabi Canada Ltd. Mesna for injection product monograph. Richmond Hill, Ontario; 30 March 2015
21. Accord Healthcare Inc. Methotrexate injection product monograph. Kirkland, Québec; October 30, 2019
22. Maia M, Farah ME, Belfort RN, et al. Effects of intravitreal triamcinolone acetonide injection with and without preservative. Br J Ophthalmol 2007;91(9):1122-1124
23. Mayne Pharma Canada. Methotrexate Product Monograph. Montreal, Quebec; December 2003
24. BC Cancer Miscellaneous Origin Tumour Group. (MOIT) BC Cancer Protocol Summary for Solid Tumours using Intrathecal Methotrexate and/or Thiotepa and/or Cytarabine. Vancouver, British Columbia: BC Cancer; September 1 2021
25. BC Cancer Lymphoma Tumour Group. (LYIT) BC Cancer Protocol Summary for Treatment of Lymphoma using Intrathecal Methotrexate and Cytarabine. Vancouver, British Columbia: BC Cancer; April 1 2022
26. Hematology/Oncology Pharmacy Association. HOPA News Clinical Pearls: Intrathecal Chemotherapy: Focus on Drugs, Dosing, and Preparation. 13(4) ed. Chicago, Illinois, USA: Hematology/Oncology Pharmacy Association; 2016
27. BC Cancer. Systemic Therapy Policy and Procedure III-50: Administration of High Alert Medications by the Intrathecal Route via Lumbar Puncture or Ommaya Reservoir. Vancouver, British Columbia; May 1 2019
28. Pfizer Canada-ULC. Methotrexate injection product monograph. Kirkland, Québec; July 8, 2019
29. Accord Healthcare Inc. Mitomycin product monograph. Kirkland, Quebec; 7 June 2017
30. Accord Healthcare Inc. Mitomycin product monograph. Kirkland, Quebec; 16 July 2018
31. Au JLS, Badalament RA, Wientjes MG, et al. Methods to improve efficacy of intravesical mitomycin C: results of a randomized phase III trial. J Natl Cancer Inst 2001;93(8):597-604
32. Jessie LS Au PharmD PhD. Professor, Ohio State University. Personal communication. 14 May 2007
33. Myers AL, Zhang Y, Kawedia JD, et al. Solubilization and stability of mitomycin C solutions prepared for intravesical administration. Drugs R D 2017;17:297-304
34. Beijnen JH, Van Gijn R, Underberg WJM. Chemical stability of the antitumor drug mitomycin C in solutions for intravesical instillation. Journal of Parenteral Science and Technology 1990;44(6):332-335
35. Teva Canada Limited. Mitomycin for injection® product monograph. Toronto, Ontario; 30 June 2017
36. Fresenius Kabi Canada Ltd. Mitoxantrone injection® product monograph. Richmond Hill, Ontario; 28 September . 2016
37. Pfizer Canada Inc. Mitoxantrone injection product monograph. Kirkland, Quebec; 17 October 2018
38. Kyowa Kirin. POTELIGEO® product monograph. Oakville, Ontario; July 12, 2023
39. Kyowa Kirin Inc. POTELIGEO® full prescribing information. Bedminster, NJ; July 2021
40. Bristol-Myers Squibb Canada Co. OPDIVO® product monograph. Montreal, Canada; February 28, 2022
41. Hoffmann-La Roche Ltd. GAZYVA® product monograph. Mississauga, Ontario; 21 December . 2015
42. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September 2007
43. Anna Sivojelezova MSc. Drug Information Associate, Hoffmann-La Roche Ltd. Personal communication. 24 April 2015
44. Omega Laboratories Limited. OCTREOTIDE ACETATE solution product monograph. Montreal, Quebec; January 25, 2022
45. Novartis Pharmaceuticals Canada Inc. SANDOSTATIN® and SANDOSTATIN® LAR® product monograph. Dorval, Quebec; April 19 2021
46. Teva Canada Limited. OCTREOTIDE for injectable suspension Product Monograph. Toronto, Ontario; Aug 30, 2021
47. Dr. Reddy's Laboratories Canada Inc. Oxaliplatin injection product monograph. Mississauga, Ontario; February 8, 2022
48. Pfizer Canada Inc. Oxaliplatin injection product monograph. Kirkland, Quebec; 31 May 2017

49. Medical Information Pfizer Canada Inc. Personal communication. 6 June 2017
50. Sandoz Canada Inc. Oxaliplatin injection product monograph. Boucherville, Quebec; 12 August 2015
51. Katryn Vosburg. Drug Information & Pharmacovigilance Specialist, Sandoz Canada Inc. Personal communication. 26 February 2016
52. Teva Canada Limited. Teva-Oxaliplatin injection® product monograph. Toronto, Ontario; 11 September 2015
53. Accord Healthcare Inc. Paclitaxel Injection. Kirkland, Quebec; September 14, 2020
54. Mercure C. Stability of 0.1 mg/mL of paclitaxel for injection in sodium chloride (0.9%) solution. St Catharines, Ontario: Biolyse Pharma; 2 February 2007
55. Biolyse. PACLITAXEL FOR INJECTION® product monograph. St. Catherines, Ontario; 12 September 2013
56. Claude Mercure. Manager, Biolyse Pharma Corporation. Personal communication. 24 June 2014
57. Zeng Z, Lazakovitch E. Study IR 120: Physical and Chemical Stability Study of Paclitaxel for Injection in 0.9 % Sodium Chloride in concentration range 0.012-0.12 mg/mL. Biolyse Pharma 2010
58. Biolyse. PACLITAXEL FOR INJECTION® product monograph. St. Catherines, Ontario; 2 December . 2005
59. Xu Q, Trissel LA, Martinez JF. Stability of paclitaxel in 5% dextrose injection or 0.9% sodium chloride injection at 4, 22, or 32 degrees C. Am J Hosp Pharm 1994;51(24):3058-60
60. Lisa Tavano. Biolyse Pharma Corporation. Personal communication. 14 May 2012
61. Sandoz Canada Inc. Paclitaxel injection USP product monograph . Boucherville, Quebec; October 18, 2021
62. Celgene Inc. ABRAXANE® product monograph. Mississauga, Ontario; 31 August 2018
63. Aisling Cahill. Drug Safety and Medical Information Specialist. Celgene Inc. Personal communication. April 23, 2015
64. Celgene Europe Limited. ABRAXANE® product monograph. Uxbridge, UK; 11 January 2013
65. Panacea Biotec Pharma Limited (distributed by Apotex Inc.). Nanoparticle, albumin-bound paclitaxel product monograph. Toronto, Ontario; November 16, 2021
66. Panacea Biotec Pharma Limited (R&D, LALRU). In-use infusion stability study report for paclitaxel powder for injectable suspension nanoparticle, albumin-bound paclitaxel (100 mg paclitaxel/vial). Toronto, Ontario; October 4, 2023
67. Pharmaceutical Partners of Canada. Pamidronate Disodium For Injection product monograph. Richmond Hill, Ontario; 18 January . 2010
68. Mayne Pharma (Canada) Inc. Pamidronate Package Insert. Montreal, Quebec; 2002
69. Omega Laboratories Ltd. Pamidronate Disodium product monograph. Montreal, Quebec; 06 June . 2005
70. Pfizer Canada-ULC. Pamidronate disodium for injection product monograph. Kirkland, Quebec; December 11, 2018
71. Sandoz Canada Inc. Pamidronate injection product monograph. Boucherville, Quebec; 28 February 2006
72. BC Cancer Agency. Pharmacy Policy Number II-20: Guiding Principles for Chemotherapy Preparation Chart. Vancouver, British Columbia: BC Cancer Agency; 19 September 2007
73. Amgen Canada. VECTIBIX® product monograph. Mississauga, Ontario; 31 March 2017
74. Amgen Inc. VECTIBIX® full prescribing information. Thousand Oaks, California USA; June 2017
75. Amgen Canada. VECTIBIX® product monograph. Mississauga, Ontario; 5 March . 2009
76. Diane Lord. Medical Information Department, Amgen Canada Inc. Personal communication. 19 June 2009
77. Servier Canada Inc. ONCASPAR® product monograph. Laval, Quebec; November 22 2021
78. Merck Canada Inc. KEYTRUDA® product monograph. Kirkland, Quebec; 6 December 2019
79. Accord Healthcare Inc. Pemetrexed disodium for injection product monograph. Kirkland, Quebec; 12 March 2015
80. Accord Healthcare Inc. Pemetrexed solution for injection product monograph. Kirkland, QC; April 26, 2021
81. Dr. Reddy's Laboratories Limited. Pemetrexed for Injection product monograph . Oakville, Ontario; April 12, 2017
82. Dr. Reddy's Laboratories Ltd. Pemetrexed for Injection product monograph. Mississauga, Ontario; May 3, 2022
83. Dr. Reddy's Laboratories Inc. Pemetrexed for injection full prescribing information (package insert). Princeton, NJ, USA; Mar 2022
84. Md Aslam. Medical information Associate. Drug information. Dr Reddy's Laboratories. Personal Communication (verbal). December 1, 2022
85. Eli Lilly and Company. ALIMTA® product monograph. Indianapolis, Indiana, USA; September 2013
86. Taro Pharmaceuticals Inc. Pemetrexed disodium for injection product monograph. Brampton, Ontario; 30 August 2018
87. Hospira Inc. NIPENT® full prescribing information. Lake Forest, Illinois USA; Oct 2019
88. Hoffmann-La Roche Limited. PERJETA® product monograph. Mississauga, Ontario; April 12, 2013
89. Hoffmann-La Roche Limited. PHESGO® product monograph. Mississauga, Ontario; January 5, 2022
90. sanofi-aventis Canada Inc. MOZOBIL® product monograph. Laval, Quebec; 8 October . 2014
91. Maureen Coughlin BSc Pharm. Solutions in Health Inc. (acting as an authorized agent of sanofi-aventis and). Personal communication. 24 May 2017
92. Hoffman-La Roche Limited. POLIVY® product monograph. Mississauga, Ontario; April 27, 2021

93. Servier Canada Inc. FOLOTYN® product monograph. Laval, Quebec; 19 October 2018
94. Marjolaine Migeon PharmD. Servier Canada Medical Information. Personal communication. 24 September 2019
95. Pfizer Canada Inc. TOMUDEX® product monograph. Kirkland, Quebec; 8 August 2017
96. Eli Lilly Canada Inc. CYRAMZA® product monograph. Toronto, Ontario; 16 July 2015
97. Marilyn Bain BScN. Senior Medical Information Associate, Eli Lilly Canada Inc. Personal communication. 16 January 2017
98. Hoffmann-La Roche Ltd. RITUXAN® product monograph. Mississauga, Ontario; October 10 2019
99. Hoffmann-La Roche Ltd. RITUXAN® SC product monograph. Mississauga, Ontario; March 21 2018
100. Sandoz Canada Inc. RIXIMYO® product monograph. Boucherville, Quebec; April 28 2020
101. Pfizer Canada-ULC. RUXIENCE® product monograph. Kirkland, Quebec; May 4 2020
102. Teva Canada Limited for Celltrion Healthcare Co Ltd. TRUXIMA® product monograph. Toronto, Ontario; July 22 2019
103. Celgene Inc. ISTODAX® product monograph. Mississauga, Ontario; 13 December 2016
104. Celgene Inc. INFO Rx ISTODAX® (romidepsin) for Injection. Mississauga, Ontario; 10 July 2017
105. Aisling Cahill. Drug Safety and Medical Information Specialist. Celgene Inc. Personal communication. 17 July 2015
106. Gilead Sciences Canada Inc. TRODELVY® product monograph. Mississauga, ON; August 2, 2022
107. Gilead Medical Information. Gilead Sciences Inc. Personal Communication: Trodelvy® (sacituzumab govitecan) Concentration After Reconstitution. Jan 24, 2022
108. Janssen Inc. SYLVANT® product monograph. Toronto, Ontario; 6 January 2016
109. Aadi Bioscience Inc. Pharmacy Manual Protocol PEX-002 (Version 4.0) Expanded Access for an Intermediate-size Population for ABI-009 (Sirolimus Albumin-bound Nanoparticles for Injectable Suspension) in Patients with Perivascular Epithelioid Cell Tumors (PEComa) or Patients with Relevant Genetic Mutations for mTOR Pathway Activation. Pacific Palisades, CA, USA; July 16, 2021
110. Aadi Bioscience Inc. ABI-009 sirolimus albumin-bound nanoparticle (packaging information - BOX). Pacific Palisades, CA; Feb 1, 2022
111. Aadi Bioscience Inc. ABI-009 sirolimus albumin-bound nanoparticle (packaging information - VIAL). Pacific Palisades, CA; Feb 1, 2022
112. Aadi Bioscience Inc. ABI-009 (*nab*-Sirolimus) Investigator's Brochure - version 10.0. Pacific Palisades, CA, USA; July 27, 2021
113. Keocyt. Streptozocin Keocyt summary of product characteristics, version 5. Montrouge, France; February 2016
114. Keocyt. ZANOSAR® package leaflet. Montrouge, France; 24 February 2016
115. Keocyt. ZANOSAR® summary of product characteristics, version 3. Montrouge, France; 31 October 2017
116. Keocyt-Riemser Pharma. Marie-Laure Vedel, Sales Order Handling and Hospital Relationship Manager. 16 September 2021
117. Medison Pharma Canada Inc. for Immunocore Ireland Limited. KIMMTRAK® product monograph. Toronto, Ontario; June 7, 2022
118. Pfeiffer, Connie. Head of Medical Affairs Immunocore. Personal Communication. Feb 16, 2022
119. Immunocore and Clinigen. MAP Pharmacy Manual – Tebentafusp 0.2 mg/mL formulation. Abingdon, Oxfordshire, UK; May 19, 2021
120. Immunocore Commercial-LLC. KIMMTRAK® full prescribing information. Conshohocken, PA, USA; Jan 2022
121. Janssen Inc. TECVAYLI® product monograph. Toronto, Ontario; July 26, 2023
122. Janssen Research & Development LLC. Investigational product preparation and Administration instructions for subcutaneous Administration of 10 mg/mL and 90 mg/mL Teclistamab (JNJ-64007957) For Weight-Based Dosing in mcg/kg for Managed Access Programs - Version 6.0. Beerse, Belgium; June 9 2023
123. de Lemos, M. BC Cancer Provincial Pharmacy Professional Practice Leader and Drug Information Coordinator. SBAR: Teclistamab Subcutaneous Injection for Multiple Myeloma at BC Cancer. (draft). BC Cancer - Systemic Therapy Program 2023
124. Pfizer Canada Inc (for Wyeth Canada). TORISEL® product monograph. Kirkland, Quebec; 21 December . 2016
125. Anna Sivojelezova M.Sc. Medical Information Associate, Wyeth. Personal communication. 6 January 2010
126. Bristol-Myers Squibb Canada. VUMON® product monograph. St. Laurent, Quebec; 26 October . 2004
127. Trissel's® 2 IV Compatibility (database on the Internet). Teniposide. Thomson Reuters MICROMEDEX® 2.0, updated periodically. Available at: <http://www.micromedex.com>. Accessed 27 April, 2011
128. Adienne-SA. TEPADINA® product monograph. Lugano, Switzerland; 28 March 2017
129. AHFS Drug Information® (database on the Internet). Thiotepa. Lexi-Comp Inc., 2018. Available at: <http://online.lexi.com>. Accessed 21 August, 2018
130. BC Cancer Miscellaneous Origin Tumour Group. (MOIT) BC Cancer Protocol Summary for Solid Tumours using Intrathecal Methotrexate and/or Thiotepa and/or Cytarabine. Vancouver, British Columbia: BC Cancer; 1 October 2018
131. Sanofi Genzyme. THYROGEN® product monograph. Mississauga, Ontario; July 29, 2021
132. Lillian Phan. Senior Manager, Medical Affairs US, EU & New Markets. Personal communication. 15 January 2021

133. BeiGene Ltd. Clinical Study Pharmacy Manual Protocol BGB-A317-207: An phase 2, open-label study of BGB-A317 in patients with relapsed or refractory mature T- and NK-cell neoplasms. (Version 5.0). San Mateo, California, USA; 8 November 2019
134. BeiGene Ltd. Tislelizumab (BGB-A317) Product Information for Investigator Sponsored Research (ISR) Investigators (Version 0.0). San Mateo, California, USA; 19 May 2020
135. Hoffmann-La Roche Limited. ACTEMRA® product monograph. Mississauga, Ontario; 27 October 2017
136. Accord Healthcare Inc. Topotecan hydrochloride for injection product monograph. Kirkland, Quebec; 9 May 2019
137. Pfizer Canada Inc. Topotecan hydrochloride for injection product monograph. Kirkland, Quebec; 5 January 2018
138. Sandoz Canada Inc. Topotecan injection product monograph. Boucherville, Quebec; 5 September . 2014
139. Hoffmann-La Roche Limited. HERCEPTIN® product monograph. Mississauga, Ontario; May 7, 2020
140. Celltrion Healthcare Co Ltd (distributed by Teva Canada Limited). HERZUMA® product monograph. Toronto, Ontario; 20 October 2020
141. BGP Pharmacy ULC. OGIVRI® product monograph. Etobicoke, Ontario; 3 May 2019
142. ULC Pfizer Canada. TRAZIMERA® product monograph. Kirkland, Quebec; 15 August 2019
143. AstraZeneca Canada Inc. ENHERTU® product monograph. Mississauga, Ontario; 15 April 2021
144. AstraZeneca Canada Inc. Medical Information. 5 October 2021
145. Hoffmann-La Roche Limited. KADCYLA® product monograph. Mississauga, Ontario; 11 September 2013
146. Hoffmann-La Roche Limited. KADCYLA® product monograph. Mississauga, Ontario; July 3, 2020
147. Medexus Inc. TRECONDYV® product monograph. Bolton, Ontario; June 25, 2021
148. Erin Wallace. Ontario Territory Manager, Medexus Pharmaceuticals Inc. Personal Communication - TRECONDYV®. November 2, 2021
149. Medical Information team. Medexus Pharmaceuticals Inc. Personal Communication - TRECONDYV®. November 1, 2021
150. medac GmbH. TRECONDI® summary of product characteristics. Wedel, Germany;
151. medac-UK. TREOSULFAN injection® summary of product characteristics. Wedel, Germany; 26 Jan 2017
152. medac-UK. TREOSULFAN injection® Details about Handling and Stability. Hamburg, Germany; August 2008
153. Henrik Fenger. Management Associate, International Division medac. Personal communication. 03 March 2010
154. medac-UK. TREOSULFAN injection® summary of product characteristics. Hamburg, Germany; 24 Jun 2008
155. ULC Pfizer Canada. Vinblastine sulfate injection product monograph. Kirkland, Quebec; 18 April 2019
156. Lexi-Drugs® (database on the Internet). VinBLASTine. Lexi-Comp Inc., 2020. Available at: <http://online.lexi.com>. Accessed 30 January, 2020
157. BC Cancer Provincial Systemic Therapy Program. Policy V-40: Dispensing and Labelling of Vinca Alkaloid Preparations. Vancouver, British Columbia: BC Cancer; 1 April 2015
158. World Health Organization. Information Exchange System - Vincristine (and other vinca alkaloids) should only be given intravenously via a minibag. Alert No. 115 ed. Geneva, Switzerland: World Health Organization; 18 July 2007
159. Teva Canada Limited. Vinblastine sulfate injection product monograph. Toronto, Ontario; 22 October 2019
160. Pfizer Canada-ULC. Vincristine sulfate injection product monograph. Kirkland, Québec; July 26, 2021
161. Teva Canada Limited. Vincristine sulfate injection® product monograph. Scarborough, Ontario; March 27, 2014
162. Pharmaceutical Partners of Canada. Vinorelbine Injection product monograph. Richmond Hill, Ontario; 15 January . 2008
163. Generic Medical Partners Inc. Vinorelbine Injection product monograph. Toronto, Ontario; 3 September 2014
164. Teva Canada Limited. Vinorelbine tartrate for Injection product monograph. Toronto, Ontario; 20 March 2014
165. Innomar Strategies Inc. (for Dr. Reddy's Laboratories Limited). Zoledronic acid for injection concentrate® product monograph. Oakville, Ontario; 11 March 2015
166. Marcan Pharmaceuticals Inc. Zoledronic acid concentrate for injection product monograph. Ottawa, Ontario; 5 February 2018
167. MDA Inc. Zoledronic acid for injection product monograph. Mississauga, Ontario; 11 August 2015
168. Novartis Pharmaceuticals Canada Inc. ZOMETA® product monograph. Dorval, Quebec; 26 July 2013
169. Sandoz Canada Inc. Zoledronic Acid - Z® product monograph. Boucherville, Quebec; 02 December 2016
170. The United States Pharmacopeia, (USP). General Chapter 797: Pharmaceutical compounding - sterile preparations. USP 27-NF 22. Rockville, Maryland: The United States Pharmacopeial Convention, Inc.; 2004
171. Kastango ES. The ASHP discussion guide for compounding sterile preparations. Bethesda (MD): American Society of Health-System Pharmacists, Inc.; 2004. p. 5